
The Stellar Imager (SI) Mission Concept

Imaging the Surfaces of Distant Stars



K. G. Carpenter (NASA/GSFC), C. J. Schrijver (LMMS) and the SI Mission Concept Development Team

Executive Summary: 23 July 2004

Mission Concept Development Team

- NASA/GSFC leads large Vision Mission Concept Development Team
 - NASA-GSFC: K. Carpenter (Study Lead), C. Bowers, W. Danchi, J. Leitner, A. Liu,

R. Lyon, L. Mazzuca, R. Moe

- LMMS/ATC: Carolus Schrijver (Science Lead), D. Chenette
- BATC: S. Kilston, C. Noecker, R. Linfield, M. Lieber, D. Miller
- NGST: R. Polidan, B. Kirkham
- JPL: N. Murphy, M. Shao, F. Hadaegh, J. Breckinridge,

G. Blackwood

- NRL/NPOI: T. Armstrong, X. Zhang
- Seabrook Eng: D. Mozurkewich
- Sigma Space: J. Marzouk
- STScI: R. Allen
- UMD: L. Mundy
- SAO: M. Karovska, J. Phillips, P. Nisenson, S. Baliunas,

S. Korzennik, W. Soon

- SUNY: F. Walter
- CU: A. Brown, G. Harper
- Potsdam U.: Klaus Strassmeier
- U Aarhus: Jörgen Christensen-Dalsgaard
- Kiepenheuer Inst: Oscar Van der Lühe


Primary Science Goals

- Study spatial and temporal stellar magnetic activity patterns in a sample of stars covering a broad range of activity level
 - Enable improved forecasting of solar activity on time scales of days to centuries
 - Understand the impact of stellar magnetic activity on planetary climates and astrobiology
- Measure internal stellar structure and rotation
- Complete the assessment of external solar systems
 - image the central stars and determine the impact of the activity of those stars on the habitability of the surrounding planets

Value to Society: Space-Weather & Earth-Climate Forecasting

- SI's observational capabilities are essential to the development and validation of a model of the solar/stellar dynamo, which is needed to
 - understand past solar activity
 - enable forecasting of solar and heliospheric activity days to decades in advance
 - anticipate the impact of those changes on the earth's biosphere and society from
 - long-term changes which effect climate, such as Maunder minima/grand maxima, which lead to lower/higher overall global temperatures and can cause crop failures, etc.
 - short-term changes, e.g. enhanced activity/flares, which have the potential to
 - disable communication satellites
 - knock out power grids
 - increase the speed of corrosion of oil pipelines
 - place astronauts at risk from particle radiation










"Strawman" Full-SI Mission Concept

- space-based UV-optical Fizeau Interferometer
 - located near the sun-earth L2 point to enable precision formation flying
 - > 20 primary mirrors distributed over virtual spherical surface 0.5 km in diameter

Capabilities Provided

- an angular resolution of 60 & 120 micro-arcsec at 1550 & 2800 Å
- ~ 1000 pixels of resolution over the surface of nearby dwarf stars
- observations in
 - − ~10-Ångstrom UV pass bands
 - C IV (100,000 K)
 - Mg II h&k (10,000 K)
 - broadband, near-UV or optical continuum (3,000-10,000 K)
- a long-term (> 10 year) mission to study stellar activity cycles:
 - individual telescopes/central hub can be refurbished or replaced

hub and primary mirrors formation fly with \sim cm precision, mirror actuators maintain optical path lengths to within 5 nm



approximate distance to hub from center of array is 65 km

Place in NASA/ESA Strategic Roadmaps


- SI is on strategic path of NASA Origins interferometry missions
 - it is a stepping stone towards crucial technology...
 - *SI* is comparable in complexity to the *Terrestrial Planet Finder*, and it may serve as a useful technological and operational pathfinder for *Planet Imager*
 - ... while addressing science goals of 3 NASA/OSS research Themes
 - understand why the sun varies (SEC)
 - understand the origin of stars, planetary systems, and life (Origins)
 - understand the structure and evolution of stars (SEU)
 - it is complementary to the planetary imaging interferometers
 - Terrestrial Planet Finder/Darwin, and Planet Imager null the stellar light to find and image planets
 - **Stellar Imager** images the central star to study the effects of that star on the habitability of planets and the formation of life on them.

TPF/Darwin, SI, and *PI* together provide complete views of other solar systems



SI and General Astrophysics

A long-baseline interferometer in space benefits many fields of astrophysics



Active Galactic Nuclei

transition zone between BLR & NLR, origin/orientation of jets

Quasi-stellar Objects & Black Holes close-in structure, radiation from accretion processes

Supernovae close-in spatial structure

Stellar interiors
internal structure in stars outside
solar parameters

Hot Stars

hot polar winds, non-radial pulsations, envelopes and shells of Be-stars

Spectroscopic binary stars

observe companions & orbits, determine stellar properties, perform key tests of stellar evolution

Interacting Binary Stars

resolve mass-exchange, dynamical evolution/accretion, study more efficient dynamos

Forming Stars/Disk systems: accretion foot-points & magnetic field structure

Cool, Evolved Giant & Supergiant Stars, and Long-Period/Semi-Regular Variables

spatiotemporal structure of extended atmospheres/winds, shocks





Current Status of SI

- SI included in far-horizon NASA SEC Roadmap
- SI selected for further concept development by the NASA HQ 2003 Vision Mission NRA review
- Major Partnerships established with LMATC, BATC, JPL, SAO, UCO to carry out further concept/technology development
- Phase I (7 primary elements) of the Fizeau Interferometry Testbed (FIT) has begun operation to develop closed-loop optical control of a multi-element array
- Next Steps
 - Continue Architecture Trade/Feasibility Studies
 - Test/demonstrate design concepts with ground-based testbed (the FIT)
 - Gather & utilize additional community input
 - Produce book summarizing science/societal motivations for mission, technology roadmap, and most promising architecture options

For more information, see: http://hires.gsfc.nasa.gov/~si